

Indefinite integral

Mathematics – FRDIS

Mendel university in Brno



europa
european
social fund in the
czech republic



EUROPEAN UNION



MINISTRY OF EDUCATION,
YOUTH AND SPORTS



OP Education
for Competitiveness

INVESTMENTS IN EDUCATION DEVELOPMENT

Podpořeno projektem Průřezová inovace studijních programů Lesnické a dřevařské fakulty MENDELU v Brně (LDF) s ohledem na disciplíny společného základu <http://akademie.ldf.mendelu.cz/cz> (reg. č. CZ.1.07/2.2.00/28.0021) za přispění finančních prostředků EU a státního rozpočtu České republiky.

Antiderivative and indefinite integral

Definition (Antiderivative)

Let f and F be functions defined on an open interval I . If

$$F'(x) = f(x) \quad \text{for every } x \in I,$$

then the function F is called an **antiderivative** of f on the interval I .

It follows from the definition: If F is an antiderivative of f , then $F + c$ (where c is any real constant), is also an antiderivative of f . Indeed,

$$[F(x) + c]' = F'(x) + c' = f(x) + 0 = f(x).$$

This means that if there exists an antiderivative of f on the given interval, then there exist infinitely many antiderivatives of f on this interval.

Example

For example, the following functions are antiderivatives of $y = x^3$:

$$y = \frac{x^4}{4}, \quad y = \frac{x^4}{4} + 3, \quad y = \frac{x^4}{4} - 7.$$

Indeed,

$$\left(\frac{x^4}{4}\right)' = x^3, \quad \left(\frac{x^4}{4} + 3\right)' = x^3, \quad \left(\frac{x^4}{4} - 7\right)' = x^3.$$

Evidently any function of the form $y = \frac{x^4}{4} + c$, where $c \in \mathbb{R}$, is an antiderivative of $y = x^3$.

Example

For example, the following functions are antiderivatives of $y = x^3$:

$$y = \frac{x^4}{4}, \quad y = \frac{x^4}{4} + 3, \quad y = \frac{x^4}{4} - 7.$$

Indeed,

$$\left(\frac{x^4}{4}\right)' = x^3, \quad \left(\frac{x^4}{4} + 3\right)' = x^3, \quad \left(\frac{x^4}{4} - 7\right)' = x^3.$$

Evidently any function of the form $y = \frac{x^4}{4} + c$, where $c \in \mathbb{R}$, is an antiderivative of $y = x^3$.

Question: Are the functions of the form $y = \frac{x^4}{4} + c$ the only antiderivatives of $y = x^3$ or can we find another antiderivative?

The following theorem gives answer to this question.

Theorem (Uniqueness of antiderivative)

Let F and G be two antiderivatives of f on an interval I . Then there exists a constant $c \in \mathbb{R}$ such that $G(x) = F(x) + c$ for every $x \in I$.

The previous theorem says that any two antiderivatives of the same function f differ only by an additive constant.

Definition (Indefinite integral)

The set of all antiderivatives of f is denoted $\int f(x) dx$ and it is called the **indefinite integral**. We write

$$\int f(x) dx = F(x) + c,$$

where F is any antiderivative of f and c is any real constant.

The symbol \int is called the **integral sign**, the function f is called **integrand**, the constant c is called the **constant of integration**. The dx is part of integral notation and indicates the variable involved. To **integrate** f means to find all antiderivatives of f . The function which has an antiderivative (indefinite integral) is called **integrable**.

It holds:

$$\left(\int f(x) dx \right)' = f(x) \quad \text{and} \quad \int (F(x))' dx = F(x) + c$$

Theorem (Sufficient condition for integrability)

Let f be a continuous function on I . Then there exists an antiderivative of f on I .

- There exist functions which are not continuous and have antiderivatives.
- There exist functions which are continuous (hence they have antiderivatives), but the antiderivatives cannot be expressed using the elementary functions. The antiderivatives of these functions, such as

$$\int e^{-x^2} dx, \quad \int \frac{e^x}{x} dx, \\ \int \frac{\sin x}{x} dx, \quad \int \frac{\cos x}{x} dx, \quad \int \sin x^2 dx, \quad \int \cos x^2 dx$$

are called **higher transcendental functions**.

Basic Formulas for integration

$$\int 0 \, dx = c$$

$$\int 1 \, dx = x + c$$

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} + c$$

$$\int e^x \, dx = e^x + c$$

$$\int a^x \, dx = \frac{a^x}{\ln a} + c$$

$$\int \frac{1}{x} \, dx = \ln |x| + c$$

$$\int \cos x \, dx = \sin x + c$$

$$\int \sin x \, dx = -\cos x + c$$

$$\int \frac{1}{\cos^2 x} \, dx = \operatorname{tg} x + c$$

$$\int \frac{1}{\sin^2 x} \, dx = -\operatorname{cotg} x + c$$

$$\int \frac{1}{\sqrt{1-x^2}} \, dx = \arcsin x + c$$

$$\int \frac{1}{x^2+1} \, dx = \operatorname{arctg} x + c$$

- We usually write $\int dx$ instead of $\int 1 dx$.
- Sometimes we write $\int \frac{dx}{f(x)}$ instead of $\int \frac{1}{f(x)} dx$.

Basic rules for integration

Theorem (Basic rules for integration)

Let f and g be functions, which are integrable on an interval I and let $c \in \mathbb{R}$. Then the following formulas hold on I :

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$
$$\int c \cdot f(x) dx = c \int f(x) dx$$

We don't have any general rule for integration products and quotients of functions and for the composite function !!!

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

Example (Integration of power functions and polynomials)

$$① \int x^3 dx = \frac{x^4}{4} + c$$

$$② \int \sqrt{x} dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}}$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3} \sqrt{x^3} + c$$

Example (Integration of power functions and polynomials)

$$① \int x^3 dx = \frac{x^4}{4} + c$$

$$② \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3} \sqrt{x^3} + c$$

$$③ \int \frac{1}{x^4} dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3}\sqrt{x^3} + c$$

$$\textcircled{3} \int \frac{1}{x^4} dx = \int x^{-4} dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3}\sqrt{x^3} + c$$

$$\textcircled{3} \int \frac{1}{x^4} dx = \int x^{-4} dx = \frac{x^{-3}}{-3}$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3}\sqrt{x^3} + c$$

$$\textcircled{3} \int \frac{1}{x^4} dx = \int x^{-4} dx = \frac{x^{-3}}{-3} = -\frac{1}{3x^3} + c$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3}\sqrt{x^3} + c$$

$$\textcircled{3} \int \frac{1}{x^4} dx = \int x^{-4} dx = \frac{x^{-3}}{-3} = -\frac{1}{3x^3} + c$$

$$\textcircled{4} \int (x^4 + 2x^3 + x - 2) dx$$

Example (Integration of power functions and polynomials)

$$\textcircled{1} \int x^3 dx = \frac{x^4}{4} + c$$

$$\textcircled{2} \int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{x^{\frac{3}{2}}}{\frac{3}{2}} = \frac{2}{3}\sqrt{x^3} + c$$

$$\textcircled{3} \int \frac{1}{x^4} dx = \int x^{-4} dx = \frac{x^{-3}}{-3} = -\frac{1}{3x^3} + c$$

$$\textcircled{4} \int (x^4 + 2x^3 + x - 2) dx = \frac{x^5}{5} + \frac{x^4}{2} + \frac{x^2}{2} - 2x + c$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}}$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\int \frac{x + 3}{x^2} \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\int \frac{x+3}{x^2} \, dx = \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\int \frac{x+3}{x^2} \, dx = \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\begin{aligned} \int \frac{x+3}{x^2} \, dx &= \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx \\ &= \ln|x| + 3 \cdot \frac{x^{-1}}{-1} \end{aligned}$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\begin{aligned} \int \frac{x+3}{x^2} \, dx &= \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx \\ &= \ln|x| + 3 \cdot \frac{x^{-1}}{-1} = \ln|x| - \frac{3}{x} + c \end{aligned}$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\begin{aligned} \int \frac{x+3}{x^2} \, dx &= \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx \\ &= \ln|x| + 3 \cdot \frac{x^{-1}}{-1} = \ln|x| - \frac{3}{x} + c \end{aligned}$$

- 3 Decomposition of the improper rational function into a sum of a polynomial and the proper rational function. (We divide the numerator by the denominator.)

$$\int \frac{x^4}{x^2+1} \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\begin{aligned} \int \frac{x+3}{x^2} \, dx &= \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx \\ &= \ln|x| + 3 \cdot \frac{x^{-1}}{-1} = \ln|x| - \frac{3}{x} + c \end{aligned}$$

- 3 Decomposition of the improper rational function into a sum of a polynomial and the proper rational function. (We divide the numerator by the denominator.)

$$\int \frac{x^4}{x^2+1} \, dx = \int \left(x^2 - 1 + \frac{1}{x^2+1} \right) \, dx$$

Example (Basic formulas and simplification of integrand)

- 1 Multiplication of the power functions

$$\int x^2 \sqrt{x} \, dx = \int x^{\frac{5}{2}} \, dx = \frac{x^{\frac{7}{2}}}{\frac{7}{2}} = \frac{2}{7} x^3 \sqrt{x} + c$$

- 2 Decomposition of the numerator of a fraction

$$\begin{aligned} \int \frac{x+3}{x^2} \, dx &= \int \left(\frac{x}{x^2} + \frac{3}{x^2} \right) \, dx = \int \left(\frac{1}{x} + 3x^{-2} \right) \, dx \\ &= \ln|x| + 3 \cdot \frac{x^{-1}}{-1} = \ln|x| - \frac{3}{x} + c \end{aligned}$$

- 3 Decomposition of the improper rational function into a sum of a polynomial and the proper rational function. (We divide the numerator by the denominator.)

$$\int \frac{x^4}{x^2+1} \, dx = \int \left(x^2 - 1 + \frac{1}{x^2+1} \right) \, dx = \frac{x^3}{3} - x + \operatorname{arctg}x + c$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\int \operatorname{tg}^2 x \, dx$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\int \operatorname{tg}^2 x \, dx = \int \frac{\sin^2 x}{\cos^2 x} \, dx$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\int \operatorname{tg}^2 x \, dx = \int \frac{\sin^2 x}{\cos^2 x} \, dx = \int \frac{1 - \cos^2 x}{\cos^2 x} \, dx$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\int \operatorname{tg}^2 x \, dx = \int \frac{\sin^2 x}{\cos^2 x} \, dx = \int \frac{1 - \cos^2 x}{\cos^2 x} \, dx = \int \left(\frac{1}{\cos^2 x} - \frac{\cos^2 x}{\cos^2 x} \right) \, dx$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\begin{aligned} \int \operatorname{tg}^2 x \, dx &= \int \frac{\sin^2 x}{\cos^2 x} \, dx = \int \frac{1 - \cos^2 x}{\cos^2 x} \, dx = \int \left(\frac{1}{\cos^2 x} - \frac{\cos^2 x}{\cos^2 x} \right) \, dx \\ &= \int \left(\frac{1}{\cos^2 x} - 1 \right) \, dx \end{aligned}$$

Example (Using formulas for trigonometric functions)

We use mainly the following formulas:

$$\operatorname{tg} x = \frac{\sin x}{\cos x}, \quad \operatorname{cotg} x = \frac{\cos x}{\sin x}, \quad \sin^2 x + \cos^2 x = 1$$

$$\begin{aligned} \int \operatorname{tg}^2 x \, dx &= \int \frac{\sin^2 x}{\cos^2 x} \, dx = \int \frac{1 - \cos^2 x}{\cos^2 x} \, dx = \int \left(\frac{1}{\cos^2 x} - \frac{\cos^2 x}{\cos^2 x} \right) \, dx \\ &= \int \left(\frac{1}{\cos^2 x} - 1 \right) \, dx = \operatorname{tg} x - x + c \end{aligned}$$

Method of substitution

Theorem (The 1 st method of substitution, $t = \varphi(x)$)

Let $f(t)$ be a function which is continuous on I and $\varphi(x)$ be a function having continuous derivative on J . Next suppose that $\varphi(x) \in I$ for every $x \in J$. Then the function $f[\varphi(x)]\varphi'(x)$ is continuous on J and it holds on this interval:

$$\int f[\varphi(x)]\varphi'(x) dx = \int f(t) dt,$$

where we substitute $t = \varphi(x)$ on the right-hand side.

- The method can be used if the integrand is of the form “**composite function $f[\varphi(x)]$ times the derivative of the interior function $\varphi'(x)$** ”
- We write a new variable t instead of the interior function $\varphi(x)$ and we write dt instead of $\varphi'(x) dx$.
- Let F be an antiderivative of f , then we proceed as follows:

$$\int f[\varphi(x)]\varphi'(x) dx = \left| \begin{array}{l} t = \varphi(x) \\ dt = \varphi'(x) dx \end{array} \right| = \int f(t) dt = F(t) + c = F[\varphi(x)] + c$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right|$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) \, dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 \, dx \\ dx = \frac{1}{3} \, dt \end{array} \right| = \frac{1}{3} \int \sin t \, dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c$$
$$= -\frac{1}{3} \cos(3x + 2) + c$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right|$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c$$
$$= -\frac{1}{3} \cos(3x + 2) + c$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt$$
$$= \frac{t^3}{3} + c$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) \, dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 \, dx \\ dx = \frac{1}{3} \, dt \end{array} \right| = \frac{1}{3} \int \sin t \, dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x \, dx &= \int (\sin x)^2 \cdot \cos x \, dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x \, dx \end{array} \right| = \int t^2 \, dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} \, dx$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) \, dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 \, dx \\ dx = \frac{1}{3} \, dt \end{array} \right| = \frac{1}{3} \int \sin t \, dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x \, dx &= \int (\sin x)^2 \cdot \cos x \, dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x \, dx \end{array} \right| = \int t^2 \, dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} \, dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x \, dx \\ x \, dx = -\frac{1}{2} \, dt \end{array} \right|$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ = -\frac{1}{3} \cos(3x + 2) + c$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ = \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt$$

Example (The 1 st method of substitution)

$$\textcircled{1} \int \sin(3x + 2) dx = \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c$$
$$= -\frac{1}{3} \cos(3x + 2) + c$$

$$\textcircled{2} \int \sin^2 x \cos x dx = \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt$$
$$= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) \, dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 \, dx \\ dx = \frac{1}{3} \, dt \end{array} \right| = \frac{1}{3} \int \sin t \, dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x \, dx &= \int (\sin x)^2 \cdot \cos x \, dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x \, dx \end{array} \right| = \int t^2 \, dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} \, dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x \, dx \\ x \, dx = -\frac{1}{2} \, dt \end{array} \right| = -\frac{1}{2} \int e^t \, dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx = \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right|$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx = \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right| = \frac{1}{4} \int \sqrt{t} dt$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx = \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right| = \frac{1}{4} \int \sqrt{t} dt = \frac{1}{4} \int t^{\frac{1}{2}} dt$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\begin{aligned} \textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx &= \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right| = \frac{1}{4} \int \sqrt{t} dt = \frac{1}{4} \int t^{\frac{1}{2}} dt \\ &= \frac{1}{4} \cdot \frac{2}{3} t^{\frac{3}{2}} + c \end{aligned}$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\begin{aligned} \textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx &= \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right| = \frac{1}{4} \int \sqrt{t} dt = \frac{1}{4} \int t^{\frac{1}{2}} dt \\ &= \frac{1}{4} \cdot \frac{2}{3} t^{\frac{3}{2}} + c = \frac{1}{6} \sqrt{t^3} + c \end{aligned}$$

Example (The 1 st method of substitution)

$$\begin{aligned} \textcircled{1} \int \sin(3x + 2) dx &= \left| \begin{array}{l} t = 3x + 2 \\ dt = 3 dx \\ dx = \frac{1}{3} dt \end{array} \right| = \frac{1}{3} \int \sin t dt = -\frac{1}{3} \cos t + c \\ &= -\frac{1}{3} \cos(3x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \sin^2 x \cos x dx &= \int (\sin x)^2 \cdot \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int t^2 dt \\ &= \frac{t^3}{3} + c = \frac{\sin^3 x}{3} + c \end{aligned}$$

$$\textcircled{3} \int x \cdot e^{1-x^2} dx = \left| \begin{array}{l} t = 1 - x^2 \\ dt = -2x dx \\ x dx = -\frac{1}{2} dt \end{array} \right| = -\frac{1}{2} \int e^t dt = -\frac{1}{2} e^t + c = -\frac{1}{2} e^{1-x^2} + c$$

$$\begin{aligned} \textcircled{4} \int x^3 \cdot \sqrt{x^4 + 1} dx &= \left| \begin{array}{l} t = x^4 + 1 \\ dt = 4x^3 dx \\ x^3 dx = \frac{1}{4} dt \end{array} \right| = \frac{1}{4} \int \sqrt{t} dt = \frac{1}{4} \int t^{\frac{1}{2}} dt \\ &= \frac{1}{4} \cdot \frac{2}{3} t^{\frac{3}{2}} + c = \frac{1}{6} \sqrt{t^3} + c = \frac{1}{6} \sqrt{(x^4 + 1)^3} + c \end{aligned}$$

Composite function with linear interior function

Let F be an antiderivative of f on an interval I . Then for $ax + b \in I$ we have:

$$\int f(ax + b) dx = \frac{1}{a}F(ax + b) + c.$$

Composite function with linear interior function

Let F be an antiderivative of f on an interval I . Then for $ax + b \in I$ we have:

$$\int f(ax + b) dx = \frac{1}{a}F(ax + b) + c.$$

Example

$$\int \sin(5x + 1) dx$$

- Using substitution:

$$\int \sin(5x + 1) dx = \left. \begin{array}{l} t = 5x + 1 \\ dt = 5 dx \\ dx = \frac{1}{5} dt \end{array} \right| = \frac{1}{5} \int \sin t dt = -\frac{1}{5} \cos t + c =$$
$$-\frac{1}{5} \cos(5x + 1) + c$$

- Using the above formula: $ax + b = 5x + 1$

$$f(x) = \sin(x) \implies F(x) = -\cos x$$

$$f(ax + b) = \sin(5x + 1) \implies F(ax + b) = -\cos(5x + 1)$$

$$\implies \int \sin(5x + 1) dx = -\frac{1}{5} \cos(5x + 1) + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

$$\textcircled{4} \int e^{2-x} \, dx$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

$$\textcircled{4} \int e^{2-x} \, dx = -e^{2-x} + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

$$\textcircled{4} \int e^{2-x} \, dx = -e^{2-x} + c$$

$$\textcircled{5} \int \sin \frac{x}{2} \, dx$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

$$\textcircled{4} \int e^{2-x} \, dx = -e^{2-x} + c$$

$$\textcircled{5} \int \sin \frac{x}{2} \, dx = \int \sin \left(\frac{1}{2}x \right) \, dx = -2 \cos \frac{x}{2} + c$$

Example

$$\textcircled{1} \int \cos 2x \, dx = \frac{1}{2} \sin 2x + c$$

$$\textcircled{2} \int (3 - 5x)^6 \, dx = -\frac{1}{5} \cdot \frac{1}{7} (3 - 5x)^7 = -\frac{1}{35} (3 - 5x)^7 + c$$

$$\textcircled{3} \int \frac{1}{2x - 3} \, dx = \frac{1}{2} \ln |2x - 3| + c$$

$$\textcircled{4} \int e^{2-x} \, dx = -e^{2-x} + c$$

$$\textcircled{5} \int \sin \frac{x}{2} \, dx = \int \sin \left(\frac{1}{2}x \right) \, dx = -2 \cos \frac{x}{2} + c$$

The integrals can be solved using the substitutions:

$$t = 2x, \quad t = 3 - 5x, \quad t = 2x - 3, \quad t = 2 - x, \quad t = \frac{x}{2}.$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

f'/f

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$① \int \frac{2x}{x^2 - 3} dx$$

f'/f

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$① \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$② \int \frac{3x}{x^2 - 3} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx = - \int \frac{-\sin x}{\cos x} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx = - \int \frac{-\sin x}{\cos x} dx = - \ln |\cos x| + c$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx = - \int \frac{-\sin x}{\cos x} dx = - \ln |\cos x| + c$$

$$\textcircled{4} \int \operatorname{cot} x dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx = - \int \frac{-\sin x}{\cos x} dx = - \ln |\cos x| + c$$

$$\textcircled{4} \int \operatorname{cot} g x dx = \int \frac{\cos x}{\sin x} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \left| \begin{array}{l} t = f(x) \\ dt = f'(x) dx \end{array} \right| = \int \frac{1}{t} dt = \ln |t| + c = \ln |f(x)| + c$$

$$\Rightarrow \boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c}$$

Example

$$\textcircled{1} \int \frac{2x}{x^2 - 3} dx = \ln |x^2 - 3| + c$$

$$\textcircled{2} \int \frac{3x}{x^2 - 3} dx = \frac{3}{2} \int \frac{2x}{x^2 - 3} dx = \frac{3}{2} \ln |x^2 - 3| + c$$

$$\textcircled{3} \int \operatorname{tg} x dx = \int \frac{\sin x}{\cos x} dx = - \int \frac{-\sin x}{\cos x} dx = - \ln |\cos x| + c$$

$$\textcircled{4} \int \operatorname{cot} g x dx = \int \frac{\cos x}{\sin x} dx = \ln |\sin x| + c$$

Trigonometric substitution

The integral of the the form $\int R(\sin x, \cos x) dx$, where R is a rational function, can be transformed (with using special substitutions) into the integral from a rational function. Special cases:

$$\textcircled{1} \int R(\sin x) \cos x dx \Rightarrow t = \sin x$$

$$\textcircled{2} \int R(\cos x) \sin x dx \Rightarrow t = \cos x$$

Remark

- Functions of the type $R(\sin x, \cos x)$: $\frac{\sin x \cos^2 x}{\sin^2 x + \cos x}$, $\frac{\cos^3 x + 1}{\sin^2 x}$
- Functions of the type $R(\sin x)$: $\frac{\sin x + 2}{\sin^2 x}$, $\frac{2 \sin x}{\sin x - 1}$
- Functions of the type $R(\cos x)$: $\frac{\cos x}{\cos^2 x + 3}$, $\frac{\cos^2 x + \cos x}{\cos x + 1}$

Example

$$\textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx$$

Example

$$\textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx = \int \frac{\sin x}{\sin^2 x + 2} \cos x dx$$

Example

$$\textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx = \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right|$$

Example

$$\textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx = \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \frac{\sin x}{\cos^3 x} dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \frac{\sin x}{\cos^3 x} dx = \int \frac{1}{\cos^3 x} \sin x dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \frac{\sin x}{\cos^3 x} dx = \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right|$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\textcircled{2} \int \frac{\sin x}{\cos^3 x} dx = \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx = \int \frac{\sin^2 x}{\cos x + 3} \sin x dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx = \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx &= \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx \\ &= \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx &= \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx \\ &= \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = \int \frac{t^2 - 1}{t + 3} dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx &= \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx \\ &= \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = \int \frac{t^2 - 1}{t + 3} dt = \int \left(t - 3 + \frac{8}{t + 3} \right) dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx &= \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx \\ &= \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = \int \frac{t^2 - 1}{t + 3} dt = \int \left(t - 3 + \frac{8}{t + 3} \right) dt \\ &= \frac{t^2}{2} - 3t + 8 \ln |t + 3| + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{\sin x \cos x}{\sin^2 x + 2} dx &= \int \frac{\sin x}{\sin^2 x + 2} \cos x dx = \left| \begin{array}{l} t = \sin x \\ dt = \cos x dx \end{array} \right| = \int \frac{t}{t^2 + 2} dt \\ &= \frac{1}{2} \int \frac{2t}{t^2 + 2} dt = \frac{1}{2} \ln(t^2 + 2) + c = \frac{1}{2} \ln(\sin^2 x + 2) + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{\sin x}{\cos^3 x} dx &= \int \frac{1}{\cos^3 x} \sin x dx = \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = - \int \frac{1}{t^3} dt \\ &= - \int t^{-3} dt = -\frac{t^{-2}}{-2} + c = \frac{1}{2t^2} + c = \frac{1}{2 \cos^2 x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} (*) \int \frac{\sin^3 x}{\cos x + 3} dx &= \int \frac{\sin^2 x}{\cos x + 3} \sin x dx = \int \frac{1 - \cos^2 x}{\cos x + 3} \sin x dx \\ &= \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \\ \sin x dx = -dt \end{array} \right| = \int \frac{t^2 - 1}{t + 3} dt = \int \left(t - 3 + \frac{8}{t + 3} \right) dt \\ &= \frac{t^2}{2} - 3t + 8 \ln |t + 3| + c = \frac{\cos^2 x}{2} - 3 \cos x + 8 \ln |\cos x + 3| + c \end{aligned}$$

Theorem (The 2 nd method of substitution, $x = \varphi(t)$)

Let $f(x)$ be a function which is continuous on I and $\varphi(t)$ be a function having continuous and nonzero derivative on J . Next suppose that $\varphi(J) = I$. Then it holds on I :

$$\int f(x) dx = \int f[\varphi(t)]\varphi'(t) dt,$$

where we substitute $t = \varphi^{-1}(x)$ on the right-hand side and where φ^{-1} is the inverse function to φ .

- We write $\varphi(t)$ instead of x and $\varphi'(t) dt$ instead of dx . The composite function obtained in the integral on the right-hand side seems to be more complicated than the original one, but in some particular cases it can be easier to integrate this composite function.
- We proceed as follows:

$$\int f(x) dx = \left| \begin{array}{l} x = \varphi(t) \\ dx = \varphi'(t) dt \end{array} \right| = \int f[\varphi(t)]\varphi'(t) dt = F(t) + c = F[\varphi^{-1}(x)] + c,$$

where $F(t)$ is an antiderivative of $f[\varphi(t)]\varphi'(t)$.

Integration of irrational function

We can eliminate roots in the integral

$$\int R(x, \sqrt[n_1]{ax+b}, \sqrt[n_2]{ax+b}, \dots) dx,$$

where R is a rational function. We use the substitution

$$t = \sqrt[s]{ax+b}, \quad \text{where } s \text{ is the least common multiple of } n_1, n_2, \dots$$

Then:

$$t = \sqrt[s]{ax+b} \implies ax+b = t^s$$

$$x = \frac{1}{a}t^s - \frac{b}{a}$$

$$dx = \frac{1}{a}st^{s-1} dt$$

Example

$$\textcircled{1} \int \frac{x}{\sqrt{x+1}} dx$$

Example

$$\textcircled{1} \int \frac{x}{\sqrt{x+1}} dx = \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right|$$

Example

$$\textcircled{1} \int \frac{x}{\sqrt{x+1}} dx = \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3}(\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx = \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right|$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx = \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctgt} + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

Example

$$\begin{aligned} 1 \quad \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} 2 \quad \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

$$3 \quad \int \frac{\sqrt{x}}{\sqrt[3]{x}+1} dx$$

Example

$$\begin{aligned} 1 \quad \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} 2 \quad \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

$$3 \quad \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx = \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right|$$

Example

$$\begin{aligned} 1 \quad \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3}(\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} 2 \quad \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

$$3 \quad \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx = \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right| = \int \frac{t^3}{t^2+1} 6t^5 dt$$

Example

$$\begin{aligned} 1 \quad \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} 2 \quad \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

$$3 \quad \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx = \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right| = \int \frac{t^3}{t^2+1} 6t^5 dt = 6 \int \frac{t^8}{t^2+1} dt$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3} (\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctg} t + c = 4 \operatorname{arctg} \sqrt{x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx &= \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right| = \int \frac{t^3}{t^2+1} 6t^5 dt = 6 \int \frac{t^8}{t^2+1} dt \\ &= 6 \int \left(t^6 - t^4 + t^2 - 1 + \frac{1}{t^2+1} \right) dt \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3}(\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctgt} + c = 4 \operatorname{arctg}\sqrt{x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx &= \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right| = \int \frac{t^3}{t^2+1} 6t^5 dt = 6 \int \frac{t^8}{t^2+1} dt \\ &= 6 \int \left(t^6 - t^4 + t^2 - 1 + \frac{1}{t^2+1} \right) dt = 6 \left(\frac{t^7}{7} - \frac{t^5}{5} + \frac{t^3}{3} - t + \operatorname{arctgt} \right) + c \end{aligned}$$

Example

$$\begin{aligned} \textcircled{1} \int \frac{x}{\sqrt{x+1}} dx &= \left| \begin{array}{l} t = \sqrt{x+1} \Rightarrow x = t^2 - 1 \\ dx = 2t dt \end{array} \right| = \int \frac{t^2 - 1}{t} 2t dt \\ &= 2 \int (t^2 - 1) dt = 2 \left(\frac{t^3}{3} - t \right) + c = \frac{2}{3}(\sqrt{x+1})^3 - 2\sqrt{x+1} + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int \frac{2}{\sqrt{x}(x+1)} dx &= \left| \begin{array}{l} t = \sqrt{x} \Rightarrow x = t^2 \\ dx = 2t dt \end{array} \right| = \int \frac{2}{t(t^2+1)} 2t dt \\ &= 4 \int \frac{1}{t^2+1} dt = 4 \operatorname{arctgt} + c = 4 \operatorname{arctg}\sqrt{x} + c \end{aligned}$$

$$\begin{aligned} \textcircled{3} \int \frac{\sqrt{x}}{\sqrt[3]{x+1}} dx &= \left| \begin{array}{l} t = \sqrt[6]{x} \Rightarrow x = t^6 \\ dx = 6t^5 dt \end{array} \right| = \int \frac{t^3}{t^2+1} 6t^5 dt = 6 \int \frac{t^8}{t^2+1} dt \\ &= 6 \int \left(t^6 - t^4 + t^2 - 1 + \frac{1}{t^2+1} \right) dt = 6 \left(\frac{t^7}{7} - \frac{t^5}{5} + \frac{t^3}{3} - t + \operatorname{arctgt} \right) + c \\ &= \frac{6\sqrt[6]{x^7}}{7} - \frac{6\sqrt[6]{x^5}}{5} + 2\sqrt{x} - 6\sqrt[6]{x} + 6 \cdot \operatorname{arctg}\sqrt[6]{x} + c \end{aligned}$$

Integration by parts

Theorem (Integration by parts formula)

Let u and v be function having continuous derivatives on an interval I . Then the following formula holds on I :

$$\int u(x)v'(x) dx = u(x)v(x) - \int u'(x)v(x) dx.$$

- 1 Integration by parts formula follows from the rule for differentiation of the product of two functions.
- 2 When using the integration by parts formula we need:
 - to differentiate the function u ,
 - to integrate the function v' . This can be a problem!!!

Moreover, for an effective application of this formula we need the product $u'v$ (which appears in the integral on the right-hand side) to be “simpler in sense of integration” than the original uv' .

Typical integrals for using integration by parts

Let P be a polynomial.

1

$$\int P(x)e^{ax+b} dx, \quad \int P(x) \sin(ax + b) dx, \quad \int P(x) \cos(ax + b) dx$$

In these cases we differentiate the polynomial and we integrate the exponential (or trigonometric) function.

2

$$\int P(x) \ln^m(ax + b) dx$$
$$\int P(x) \operatorname{arctg}(ax + b) dx, \quad \int P(x) \operatorname{arccotg}(ax + b) dx$$
$$\int P(x) \arcsin(ax + b) dx, \quad \int P(x) \arccos(ax + b) dx$$

In these cases we integrate the polynomial and we differentiate the logarithmic or the cyclometric function. The case $P(x) = 1$ is also included.

Example (Integration by parts I)

$$\textcircled{1} \int x \cos x \, dx$$

Example (Integration by parts I)

$$\textcircled{1} \int x \cos x \, dx = \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right|$$

Example (Integration by parts I)

$$\textcircled{1} \int x \cos x \, dx = \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\textcircled{2} \int (x^2 + 1)e^{-x} \, dx$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\textcircled{2} \int (x^2 + 1)e^{-x} \, dx = \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right|$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx = \left| \begin{array}{ll} u = x & v' = e^{-x} \\ u' = 1 & v = -e^{-x} \end{array} \right| \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx = \left| \begin{array}{ll} u = x & v' = e^{-x} \\ u' = 1 & v = -e^{-x} \end{array} \right| \\ &= -e^{-x}(x^2 + 1) + 2 \left[x(-e^{-x}) - \int 1 \cdot (-e^{-x}) \, dx \right] \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx = \left| \begin{array}{ll} u = x & v' = e^{-x} \\ u' = 1 & v = -e^{-x} \end{array} \right| \\ &= -e^{-x}(x^2 + 1) + 2 \left[x(-e^{-x}) - \int 1 \cdot (-e^{-x}) \, dx \right] \\ &= -e^{-x}(x^2 + 1) - 2xe^{-x} + 2 \int e^{-x} \, dx \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx = \left| \begin{array}{ll} u = x & v' = e^{-x} \\ u' = 1 & v = -e^{-x} \end{array} \right| \\ &= -e^{-x}(x^2 + 1) + 2 \left[x(-e^{-x}) - \int 1 \cdot (-e^{-x}) \, dx \right] \\ &= -e^{-x}(x^2 + 1) - 2xe^{-x} + 2 \int e^{-x} \, dx \\ &= -e^{-x}(x^2 + 1) - 2xe^{-x} - 2e^{-x} + c \end{aligned}$$

Example (Integration by parts I)

$$\begin{aligned} \textcircled{1} \int x \cos x \, dx &= \left| \begin{array}{ll} u = x & v' = \cos x \\ u' = 1 & v = \sin x \end{array} \right| = x \sin x - \int \sin x \, dx \\ &= x \sin x + \cos x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int (x^2 + 1)e^{-x} \, dx &= \left| \begin{array}{ll} u = x^2 + 1 & v' = e^{-x} \\ u' = 2x & v = -e^{-x} \end{array} \right| \\ &= (x^2 + 1)(-e^{-x}) - \int 2x(-e^{-x}) \, dx \\ &= -e^{-x}(x^2 + 1) + 2 \int xe^{-x} \, dx = \left| \begin{array}{ll} u = x & v' = e^{-x} \\ u' = 1 & v = -e^{-x} \end{array} \right| \\ &= -e^{-x}(x^2 + 1) + 2 \left[x(-e^{-x}) - \int 1 \cdot (-e^{-x}) \, dx \right] \\ &= -e^{-x}(x^2 + 1) - 2xe^{-x} + 2 \int e^{-x} \, dx \\ &= -e^{-x}(x^2 + 1) - 2xe^{-x} - 2e^{-x} + c = -e^{-x}(x^2 - 2x + 3) + c \end{aligned}$$

Example (Integration by parts II)

① $\int \ln x \, dx$

Example (Integration by parts II)

$$\textcircled{1} \int \ln x \, dx = \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right|$$

Example (Integration by parts II)

$$\textcircled{1} \int \ln x \, dx = \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

$$\textcircled{2} \int x \operatorname{arctg} x \, dx$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

$$\textcircled{2} \int x \operatorname{arctg} x \, dx = \left| \begin{array}{ll} u = \operatorname{arctg} x & v' = x \\ u' = \frac{1}{x^2+1} & v = \frac{x^2}{2} \end{array} \right|$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

$$\textcircled{2} \int x \operatorname{arctg} x \, dx = \left| \begin{array}{ll} u = \operatorname{arctg} x & v' = x \\ u' = \frac{1}{x^2+1} & v = \frac{x^2}{2} \end{array} \right| = \frac{x^2}{2} \operatorname{arctg} x - \frac{1}{2} \int \frac{x^2}{x^2+1} \, dx$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int x \arctg x \, dx &= \left| \begin{array}{ll} u = \arctg x & v' = x \\ u' = \frac{1}{x^2+1} & v = \frac{x^2}{2} \end{array} \right| = \frac{x^2}{2} \arctg x - \frac{1}{2} \int \frac{x^2}{x^2+1} \, dx \\ &= \frac{x^2}{2} \arctg x - \frac{1}{2} \int \left(1 - \frac{1}{x^2+1} \right) dx \end{aligned}$$

Example (Integration by parts II)

$$\begin{aligned} \textcircled{1} \int \ln x \, dx &= \left| \begin{array}{ll} u = \ln x & v' = 1 \\ u' = \frac{1}{x} & v = x \end{array} \right| = x \ln x - \int \frac{1}{x} \cdot x \, dx \\ &= x \ln x - \int dx = x \ln x - x + c \end{aligned}$$

$$\begin{aligned} \textcircled{2} \int x \arctg x \, dx &= \left| \begin{array}{ll} u = \arctg x & v' = x \\ u' = \frac{1}{x^2+1} & v = \frac{x^2}{2} \end{array} \right| = \frac{x^2}{2} \arctg x - \frac{1}{2} \int \frac{x^2}{x^2+1} \, dx \\ &= \frac{x^2}{2} \arctg x - \frac{1}{2} \int \left(1 - \frac{1}{x^2+1} \right) dx = \frac{x^2}{2} \arctg x - \frac{1}{2} (x - \arctg x) + c \end{aligned}$$

Using the computer algebra systems

Wolfram Alpha:

`http://www.wolframalpha.com/`

Mathematical Assistant on Web (MAW):

`wood.mendelu.cz/math/maw-html/index.php?lang=en&form=integral`

Example

Using the Wolfram Alpha find the integral

$$\int \ln x \, dx.$$

Solution:

`integrate ln x dx`